

Double q -Analytic q -Hermite Binomial Formula and q -Traveling Waves

Sengul Nalci and Oktay K. Pashaev

Department of Mathematics, Izmir Institute of Technology
Urla-Izmir, 35430, Turkey

February 3, 2016

Abstract

Motivated by derivation of the Dirac type δ -function for quantum states in Fock-Bargmann representation, we find q -binomial expansion in terms of q -Hermite polynomials, analytic in two complex arguments. Based on this representation, we introduce a new class of complex functions of two complex arguments, which we call the double q -analytic functions. The real version of these functions describe the q -analogue of traveling waves, which is not preserving the shape during evolution as the usual traveling wave. For corresponding q -wave equation we solve IVP in the q -D'Alembert form.

1 Introduction

In complex analysis, a complex function $f(z)$ of one complex variable z is analytic in some domain D if in D it satisfies

$$\frac{\partial}{\partial \bar{z}} f(z) = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) f(z) = 0. \quad (1)$$

If we consider a complex function $f(z, w)$ of two complex variables z, w , analytic in both variables

$$\frac{\partial}{\partial \bar{z}} f(z, w) = \frac{\partial}{\partial \bar{w}} f(z, w) = 0, \quad (2)$$

it is called a double analytic if

$$\frac{1}{2} \left(\frac{\partial}{\partial z} + i \frac{\partial}{\partial w} \right) f(z, w) = 0. \quad (3)$$

As an example, $f(z, w) = (z + iw)^2$ is double analytic, while $(z - iw)^2$ is double anti-analytic.

In general, double analytic function can be written as power series in complex binomials

$$f(z, w) = \sum_{n=0}^{\infty} a_n (z + iw)^n.$$

For such double analytic binomials here we derive the following Hermite binomial formula

$$(z + iw)^n = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} i^k H_{n-k}(z) H_k(w). \quad (4)$$

Derivation of this formula is motivated by description of the Dirac type δ -function for quantum states in the Fock-Bargmann representation. As is well known, states of a quantum system in the Fock-Bargmann representation are described by complex analytic function $f(z)$ and visa versa [1]. In this representation, due to formula

$$\int d\mu(z) e^{\xi \bar{z}} f(z) = f(\xi), \quad (5)$$

where measure $d\mu(z) = dz d\bar{z} e^{-z\bar{z}}$, the exponential function plays the role of the Dirac type δ function [2]. Proof of this formula is based on following identity

$$\int d\mu(z) e^{\xi \bar{z}} z^n = \xi^n. \quad (6)$$

By generalization of this identity to two complex variables we find expansion (4).

In the present paper, we are going not only proof the identity (4), but also derive the q -analogue of this identity.

Recently, [5] we have introduced a complex function $f(z; q)$ of one complex variable z according to equation

$$D_q^{\bar{z}} f(z; q) = \frac{1}{2} \left(D_q^x + i D_q^y \right) f(z; q) = 0 \quad (7)$$

and we called it as the q -analytic function. We have described a wide class of these functions, which are not analytic in the usual sense (1).

Example: Complex q -binomial

$$(x + iy)_q^2 = (x + iy)(x + iqy) = \frac{1}{2} ((1 + q)z^2 + (1 - q)\bar{z}z)$$

is not analytic $\frac{\partial}{\partial \bar{z}}(x + iy)_q^2 \neq 0$, but q -analytic $D_q^{\bar{z}}(x + iy)_q^2 = 0$.

A complex function $f(z, w)$ of two complex variables z and w , analytic in both variables and satisfying equation

$$\bar{D}_{z,w} f(z, w) = \frac{1}{2} \left(D_q^z + i D_q^w \right) f(z, w) = 0 \quad (8)$$

we call the *double q -analytic* function.

Example: Complex q -binomial

$$(z + iw)_q^2 = z^2 + [2]_q iwz - qw^2$$

is analytic in z and w , since $\frac{\partial}{\partial \bar{z}}(z + iw)_q^2 = \frac{\partial}{\partial \bar{w}}(z + iw)_q^2 = 0$ and double q -analytic due to $\bar{D}_{z,w}(z + iw)_q^2 = 0$.

We will show that complex q -binomial $(z + iw)_q^n$, for n -positive integer, is double q -analytic. This is why any convergent power series

$$f(z, w)_q = \sum_{n=0}^{\infty} a_n (z + iw)_q^n$$

represents a double q -analytic function. As a central result of the paper, we find an expansion of the q -binomial in terms of q -Hermite polynomials

$$(z + iw)_q^n = \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q i^k q^{\frac{k(k-1)}{2}} H_{n-k}(z; q) H_k(qw, \frac{1}{q}). \quad (9)$$

In the limit $q \rightarrow 1$, this formula reduces to the Hermite binomial formula (4).

As an application of our results, we consider q -traveling waves in the form of power series of real q -binomial $(x \pm ct)_q^n$. For these traveling waves we find q -Hermite binomial expansion in terms of x and t variables. We notice that in contrast to usual traveling waves, the q -traveling waves are not preserving the shape during evolution. The q -traveling waves are subject to q -wave equation

$$\left((D_{\frac{1}{q}}^t)^2 - c^2 (D_q^x)^2 \right) u(x, t) = 0,$$

for which we solve IVP in the q -D'Alembert form with the Jackson integral representation.

The paper organized as follows. In Section 2 we derive the Hermite binomial formula (4). In Section 3 we introduce q -Hermite polynomials and derive the q -Hermite binomial formula. The q -binomials are the double q -analytic functions, as we demonstrate in Section 4. In Section 5 we discuss relation between the double q -analytic binomial and the q -analytic one. In Section 6 we introduce q -traveling waves in terms of real q -binomials and solve IVP for q -wave equation in D'Alembert form. We illustrate our results for different initial values by particular examples in Section 7. Finally, in Section 8, we derive q -Hermite polynomial expansion of q -traveling waves.

2 Hermite Binomial Formula

In this section, we derive the Hermite binomial formula (4). We start with the following Lemma:

Lemma 2.0.1 For arbitrary complex numbers ξ and η :

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{d^{n-k}}{d\xi^{n-k}} \frac{d^k}{d\eta^k} e^{\xi^2/4 - \eta^2/4} = \left(\frac{\xi + \eta}{2} \right)^n e^{\xi^2/4 - \eta^2/4} \quad (10)$$

Proof 2.0.2

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{d^{n-k}}{d\xi^{n-k}} \frac{d^k}{d\eta^k} e^{\xi^2/4 - \eta^2/4} = \left(\frac{d}{d\xi} - \frac{d}{d\eta} \right)^n e^{\xi^2/4 - \eta^2/4} \quad (11)$$

By changing variables ξ and η to λ and μ according to

$$\lambda + \mu = \xi, \quad \lambda - \mu = \eta,$$

we have

$$\lambda \mu = \frac{\xi^2 - \eta^2}{4} = \left(\frac{\xi - \eta}{2} \right) \left(\frac{\xi + \eta}{2} \right)$$

and

$$\left(\frac{d}{d\xi} - \frac{d}{d\eta} \right)^n e^{\lambda \mu} = \left(\frac{d}{d\mu} \right)^n e^{\lambda \mu} = \lambda^n e^{\lambda \mu} = \left(\frac{\xi + \eta}{2} \right)^n e^{\xi^2/4 - \eta^2/4}. \quad \diamond \quad (12)$$

Corollary 2.0.3

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \left(\frac{d^{n-k}}{d\xi^{n-k}} e^{\xi^2/4} \right) \left(\frac{d^k}{d\xi^k} e^{-\xi^2/4} \right) = \xi^n \quad (13)$$

Proof 2.0.4 By taking the limit of (10) as $\eta \rightarrow \xi \Rightarrow \mu \rightarrow 0$ and $\lambda \rightarrow \xi$,

$$\lim_{\eta \rightarrow \xi} \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{d^{n-k}}{d\xi^{n-k}} \frac{d^k}{d\eta^k} e^{\xi^2/4 - \eta^2/4} = \lim_{\mu \rightarrow 0, \lambda \rightarrow \xi} \lambda^n e^{\lambda \mu} = \xi^n. \quad (14)$$

Lemma 2.0.5 For an arbitrary complex number ξ and $n = 1, 2, \dots$,

$$\int dz d\bar{z} e^{-z\bar{z}} e^{\xi\bar{z}} z^n = \xi^n. \quad (15)$$

Proof 2.0.6 By changing complex coordinates to the Cartesian ones, the integral is expressed in terms of the sum

$$\begin{aligned} \int dz d\bar{z} e^{-z\bar{z}} e^{\xi\bar{z}} z^n &= \frac{1}{\pi} \int dx dy e^{\xi(x-iy)} e^{-(x^2+y^2)} (x+iy)^n \\ &= \frac{1}{\pi} \sum_{k=0}^n \binom{n}{k} i^k \int dx x^{n-k} e^{-x^2} e^{\xi x} \int dy y^k e^{-y^2} e^{-i\xi y} \\ &= \frac{1}{\pi} \sum_{k=0}^n \binom{n}{k} i^k \frac{d^{n-k}}{d\xi^{n-k}} \int dx e^{-x^2 + \xi x} \frac{d^k}{d(-i\xi)^k} \int dy e^{-y^2 - i\xi y} \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{d^{n-k}}{d\xi^{n-k}} e^{\xi^2/4} \frac{d^k}{d\xi^k} e^{-\xi^2/4}, \end{aligned} \quad (16)$$

where we have used the Gaussian integrals

$$\int_{-\infty}^{\infty} e^{-x^2+ax} dx = \sqrt{\pi} e^{a^2/4}$$

and

$$\int_{-\infty}^{\infty} e^{-x^2+ibx} dx = \sqrt{\pi} e^{-b^2/4}.$$

By using Corollary 2.0.3. we find desired result (15).

We can generalize this result for an arbitrary analytic function given by power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$, so that,

$$\int d\mu(z) e^{\xi \bar{z}} f(z) = f(\xi).$$

The above proof implies some interesting binomial identity for Hermite polynomials. We start from the Rodrigues formula for Hermite polynomials:

$$H_n(z) = (-1)^n e^{z^2} \frac{d^n}{dz^n} e^{-z^2} \quad (17)$$

and by replacing $z \rightarrow iz$

$$H_n(iz) = i^n e^{-z^2} \frac{d^n}{dz^n} e^{z^2}. \quad (18)$$

Then we have the following

Identity 2.0.7

$$\frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} (-i)^{n-k} H_{n-k}\left(\frac{i}{2}\xi\right) H_k\left(\frac{\xi}{2}\right) = \xi^n. \quad (19)$$

Proof 2.0.8 According to previous proof

$$\xi^n = \int dz d\bar{z} e^{-z\bar{z}} e^{\xi \bar{z}} z^n = \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{d^{n-k}}{d\xi^{n-k}} e^{\xi^2/4} \frac{d^k}{d\xi^k} e^{-\xi^2/4}. \quad (20)$$

By inserting $1 = e^{\xi^2/4} e^{-\xi^2/4}$ and using Rodrigues formulas we have

$$\begin{aligned} & \sum_{k=0}^n \binom{n}{k} (-1)^k e^{\xi^2/4} \left(e^{-\xi^2/4} \frac{d^{n-k}}{d\xi^{n-k}} e^{\xi^2/4} \right) \left(\frac{d^k}{d\xi^k} e^{-\xi^2/4} \right) \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^k \left(\frac{1}{2i} \right)^{n-k} H_{n-k}\left(\frac{i}{2}\xi\right) \left(e^{\xi^2/4} \frac{d^k}{d\xi^k} e^{-\xi^2/4} \right) \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^k \left(\frac{1}{2i} \right)^{n-k} H_{n-k}\left(i\frac{\xi}{2}\right) \left(-\frac{1}{2} \right)^k H_k\left(\frac{\xi}{2}\right) \\ &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} i^{n-k} H_{n-k}\left(i\frac{\xi}{2}\right) H_k\left(\frac{\xi}{2}\right) \\ &= \xi^n \quad \diamond \end{aligned} \quad (21)$$

Particular forms of this identity are:

$$\frac{1}{2^{2n}} \sum_{k=0}^n \binom{n}{k} i^k H_{n-k}(z) H_k(-iz) = z^n, \quad (\xi \rightarrow -2iz) \quad (22)$$

and (by reductions $\xi = x$ and $\xi = iy$),

$$\frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} i^{n-k} H_{n-k}\left(\frac{ix}{2}\right) H_k\left(\frac{x}{2}\right) = x^n, \quad (23)$$

$$\frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} i^{n-k} H_{n-k}\left(\frac{-y}{2}\right) H_k\left(\frac{y}{2}\right) = i^n y^n. \quad (24)$$

This result can be generalized to the Hermite binomial formula with two complex variables z and w

Identity 2.0.9

$$(z + iw)^n = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} i^k H_{n-k}(z) H_k(w) \quad (25)$$

Proof 2.0.10 *From generating function for Hermite polynomials*

$$g(z, t) = e^{-t^2 + 2tz} = \sum_{n=0}^{\infty} H_n(z) \frac{t^n}{n!} \quad (26)$$

and

$$g(w, \tau) = e^{-\tau^2 + 2\tau w} = \sum_{k=0}^{\infty} H_k(w) \frac{\tau^k}{k!} \quad (27)$$

by changing variable $\tau = it$ in the second one and multiplying (26) and (27) we have

$$g(z, t) g(w, it) = e^{2t(z+iw)} = \sum_{l=0}^{\infty} \sum_{k=0}^{\infty} \frac{H_l(z)}{l!} \frac{H_k(w)}{k!} i^k t^{l+k}. \quad (28)$$

By changing the order of double sum with $l+k = n$, and expanding the left hand side in t we get

$$g(z, t) g(w, it) = \sum_{n=0}^{\infty} \frac{2^n t^n (z + iw)^n}{n!} = \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{k=0}^n \binom{n}{k} H_{n-k}(z) H_k(w) i^k. \quad (29)$$

By equating terms of the same power t^n we obtain the desired result (25):

$$(z + iw)^n = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} i^k H_{n-k}(z) H_k(w). \quad \diamond$$

Here we can give also another proof of this relation by using the holomorphic Laplace equation.

Proof 2.0.11 $\zeta^n \equiv (z + iw)^n$ is double analytic function of two complex variables z and w . Therefore, due to (3) it satisfies the holomorphic Laplace equation $\Delta \zeta^n = 0$, where $\Delta \equiv \frac{\partial^2}{\partial z^2} + \frac{\partial^2}{\partial w^2} = \left(\frac{\partial}{\partial z} - \frac{\partial}{\partial w}\right) \left(\frac{\partial}{\partial z} + \frac{\partial}{\partial w}\right)$, which implies $\Delta^k \zeta^n = 0$ for every $k = 1, 2, \dots$ Expanding

$$e^{-\frac{1}{4}\Delta}(z+iw)^n = \left(1 - \frac{1}{4}\Delta + \frac{(-\frac{1}{4}\Delta)^2}{2!} + \dots + \frac{(-\frac{1}{4}\Delta)^n}{n!} + \dots\right)(z+iw)^n = (z+iw)^n \quad (30)$$

we have

$$\begin{aligned} e^{-\frac{1}{4}\Delta}(z+iw)^n &= e^{-\frac{1}{4}\frac{d^2}{dz^2}} e^{-\frac{1}{4}\frac{d^2}{dw^2}} \sum_{k=0}^n \binom{n}{k} z^{n-k} i^k w^k \\ &= \sum_{k=0}^n \binom{n}{k} i^k \left(e^{-\frac{1}{4}\frac{d^2}{dz^2}} z^{n-k}\right) \left(e^{-\frac{1}{4}\frac{d^2}{dw^2}} w^k\right). \end{aligned} \quad (31)$$

Then due to relation:

$$H_n(z) = 2^n e^{-\frac{1}{4}\frac{d^2}{dz^2}} z^n$$

we get the Hermit binomial formula

$$(z+iw)^n = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} H_{n-k}(z) H_k(w) i^k. \quad (32)$$

3 q-Hermite Binomial Formula

Here we are going to generalize the above formula to the q -binomial case. For this, first we need to introduce the q -Hermite polynomials.

3.1 q -Hermite Polynomials

In paper [4], studying the q -heat and q -Burgers equations, we have defined the q -Hermite polynomials according to generating function

$$e_q(-t^2)e_q([2]_q tx) = \sum_{n=0}^{\infty} H_n(x; q) \frac{t^n}{[n]_q!}, \quad (33)$$

where

$$e_q(x) = \sum_{n=0}^{\infty} \frac{x^n}{[n]_q!}, \quad E_q(x) = \sum_{n=0}^{\infty} q^{\frac{n(n-1)}{2}} \frac{x^n}{[n]_q!}$$

are Jackson's q -exponential functions and q -numbers and q -factorials are defined as follows:

$$[n]_q = \frac{q^n - 1}{q - 1}, \quad [n]_q! = [1]_q [2]_q \dots [n]_q.$$

By q -differentiating the generating function (33) according to x and t we have the recurrence relations, correspondingly

$$D_x H_n(x; q) = [2]_q [n]_q H_{n-1}(x; q), \quad (34)$$

$$H_{n+1}(x; q) = [2]_q x H_n(x; q) - [n]_q H_{n-1}(qx; q) - [n]_q q^{\frac{n+1}{2}} H_{n-1}(\sqrt{q}x; q). \quad (35)$$

We get also the special values

$$H_{2n}(0; q) = (-1)^n \frac{[2n]_q!}{[n]_q!}, \quad (36)$$

$$H_{2n+1}(0; q) = 0, \quad (37)$$

and the parity relations

$$H_n(-x; q) = (-1)^n H_n(x; q). \quad (38)$$

For more details we refer to paper [4]. First few polynomials are

$$H_0(x; q) = 1, \quad H_1(x; q) = [2]_q x,$$

$$H_2(x; q) = [2]_q^2 x^2 - [2]_q, \quad H_3(x; q) = [2]_q^3 x^3 - [2]_q^2 [3]_q x,$$

$$H_4(x; q) = [2]_q^4 x^4 - [2]_q^2 [3]_q [4]_q x^2 + [2]_q [3]_q [2]_q^2,$$

and when $q \rightarrow 1$ these polynomials reduce to the standard Hermite polynomials.

The generating function (33) for $t = 1$ gives expansion of q -exponential function in terms of q -Hermite polynomials

$$\sum_{n=0}^{\infty} \frac{H_n(x; q)}{[n]_q!} = \frac{e_q([2]_q x)}{E_q(1)}. \quad (39)$$

In the limiting case $q \rightarrow 1$ it gives expansion of exponential

$$e^x = e \sum_{n=0}^{\infty} \frac{H_n\left(\frac{x}{2}\right)}{n!}$$

and for $x = 1$ we find for Euler's number e as:

$$e = \sum_{n=0}^{\infty} \frac{H_n(1)}{n!}.$$

For the Jackson q -exponential function we have the q -analog of this expansion:

$$e_q(x) = e_{\frac{1}{q}} \sum_{n=0}^{\infty} \frac{H_n\left(\frac{x}{[2]_q}; q\right)}{[n]_q!},$$

and as follows

$$\frac{e_q}{e_{\frac{1}{q}}} = e_q(1)e_q(-1) = \sum_{n=0}^{\infty} \frac{H_n\left(\frac{1}{[2]_q}; q\right)}{[n]_q!}, \quad (40)$$

where

$$e_q = \sum_{n=0}^{\infty} \frac{1}{[n]_q!}$$

is the q -analog of Euler's number e . Relation (40) should be compared with the next one

$$e_q(1)e_q(-1) = e_{q^2} \left(\frac{1-q}{1+q} \right) = \prod_{k=1}^{\infty} \frac{1}{1 - (1-q^2)q^k}$$

which is coming for $x = 1$ from the following identity.

Identity 3.1.1

$$e_q(x)e_q(-x) = e_{q^2} \left(\frac{1-q}{1+q} x^2 \right).$$

Proof 3.1.2 *We expand and change order of the double sum*

$$e_q(x)e_q(-x) = \sum_{k=0}^{\infty} \frac{x^k}{[k]_q!} \sum_{l=0}^{\infty} \frac{(-1)^l x^l}{[l]_q!} = \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} (-1)^l \frac{x^{k+l}}{[k]_q! [l]_q!} = \quad (41)$$

$$\sum_{n=0}^{\infty} \frac{x^n}{[n]_q!} \sum_{k=0}^n (-1)^k \frac{[n]_q!}{[k]_q! [n-k]_q!} = \sum_{n=0}^{\infty} \frac{x^n}{[n]_q!} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (-1)^k. \quad (42)$$

Splitting the first sum to the even $n = 2m$ and to the odd parts $n = 2m + 1$ we have

$$\sum_{m=0}^{\infty} \frac{x^{2m}}{[2m]_q!} \sum_{k=0}^{2m} \begin{bmatrix} 2m \\ k \end{bmatrix}_q (-1)^k + \sum_{m=0}^{\infty} \frac{x^{2m+1}}{[2m+1]_q!} \sum_{k=0}^{2m+1} \begin{bmatrix} 2m+1 \\ k \end{bmatrix}_q (-1)^k. \quad (43)$$

Due to known identities [3]

$$\sum_{k=0}^{2m+1} \begin{bmatrix} 2m+1 \\ k \end{bmatrix}_q (-1)^k = 0,$$

$$\sum_{k=0}^{2m} \begin{bmatrix} 2m \\ k \end{bmatrix}_q (-1)^k = (1-q)(1-q^3)\dots(1-q^{2m-3})(1-q^{2m-1}),$$

the second sum vanishes and for the first sum we get

$$\sum_{m=0}^{\infty} \frac{x^{2m}}{[2m]_q!} (1-q)(1-q^3)\dots(1-q^{2m-3}).$$

Since $[2m]_q = [2]_q [m]_{q^2}$, we can rewrite this sum as

$$\sum_{m=0}^{\infty} \frac{x^{2m}(1-q)^m}{[2]_q^m [m]_{q^2}!} = e_{q^2} \left(\frac{(1-q)x^2}{[2]_q} \right).$$

Finally,

$$e_q(x)e_q(-x) = e_{q^2} \left(\frac{1-q}{1+q} x^2 \right).$$

3.2 q -Hermite Binomials

Here we formulate our main result as the q -Hermite binomial identity.

Identity 3.2.1 *The q -analogue of identity (25), giving q -binomial expansion in terms of q -Hermite polynomials is*

$$(z + iw)_q^n = \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q i^k q^{\frac{k(k-1)}{2}} H_{n-k}(z; q) H_k(qw, \frac{1}{q}) \quad (44)$$

Proof 3.2.2 *By using the generating function for q -Hermite polynomials (33) and replacing $x \rightarrow Z$ we obtain*

$$e_q(-t^2)e_q([2]_q Zt) = \sum_{n=0}^{\infty} H_n(Z; q) \frac{t^n}{[n]_q!}. \quad (45)$$

In this formula we replace $t \rightarrow it$, $Z \rightarrow W$, $q \rightarrow 1/q$ so that

$$e_{\frac{1}{q}}(t^2)e_{\frac{1}{q}}([2]_{\frac{1}{q}} iWt) = \sum_{n=0}^{\infty} H_n(W; \frac{1}{q}) i^n \frac{t^n}{[n]_{\frac{1}{q}}!}. \quad (46)$$

Multiplying (45) with (46) and using factorization of q -exponential functions

$$e_q(x)e_{\frac{1}{q}}(y) = \sum_{n=0}^{\infty} \frac{(x+y)_q^n}{[n]_q!} \equiv e_q(x+y)_q \quad (47)$$

leading to

$$e_q(-t^2)e_{\frac{1}{q}}(t^2) = e_q(0)_q = 1,$$

we find

$$e_q(t([2]_q Z + [2]_{\frac{1}{q}} iW))_q = \sum_{l=0}^{\infty} \sum_{k=0}^{\infty} \frac{H_l(Z; q)}{[l]_q!} \frac{H_k(W; \frac{1}{q}) i^k}{[k]_{\frac{1}{q}}!} t^{l+k}. \quad (48)$$

By changing order of the double sum and expanding the left hand side in t , we get

$$\sum_{n=0}^{\infty} \frac{t^n ([2]_q Z + [2]_{\frac{1}{q}} iW)_q^n}{[n]_q!} = \sum_{n=0}^{\infty} \frac{t^n}{[n]_q!} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} H_{n-k}(Z; q) H_k(W; \frac{1}{q}) i^k. \quad (49)$$

Then, at power t^n we have identity

$$\left(Z + i\frac{W}{q}\right)_q^n = \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} H_{n-k}(Z; q) H_k(W; \frac{1}{q}) i^k, \quad (50)$$

where

$$[k]_{\frac{1}{q}} = \frac{1}{q^{k-1}} [k]_q, \quad [k]_{\frac{1}{q}}! = \frac{1}{q^{\frac{k(k-1)}{2}}} [k]_q!$$

By replacing $Z = z$ and $\frac{W}{q} = w$ the desired result is obtained

$$(z + iw)_q^n = \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} H_{n-k}(z; q) H_k(qw; \frac{1}{q}) i^k. \quad \diamond$$

4 Double q-Analytic Function

Here we consider a class of complex valued functions of two complex variables, z and w , (or four real variables), analytic in these variables $\frac{\partial}{\partial \bar{z}} f = \frac{\partial}{\partial \bar{w}} f = 0$.

Definition 4.0.3 A complex-valued function $f(z, w)$ of four real variables is called the double analytic in a region if the following identity holds in the region:

$$\bar{\partial}_{z,w} f \equiv \frac{1}{2}(\partial_z + i\partial_w) f = 0, \quad (51)$$

where

$$\partial_z f = \frac{1}{2}(\partial_x - i\partial_y), \quad \partial_w f = \frac{1}{2}(\partial_u - i\partial_v)$$

and $z = x + iy$, $w = u + iv$.

Definition 4.0.4 A complex-valued function $f(z, w)$ of four real variables is called the double q -analytic in a region if the following identity holds in the region:

$$\bar{D}_{z,w} f = \frac{1}{2}(D_q^z + iD_{\frac{1}{q}}^w) f = 0, \quad (52)$$

where

$$D_q^z f(z, w) = \frac{f(qz, w) - f(z, w)}{(q-1)z}, \quad D_{\frac{1}{q}}^w f(z, w) = \frac{f(z, \frac{w}{q}) - f(z, w)}{(\frac{1}{q}-1)w}$$

and $z = x + iy$, $w = u + iv$.

Here we should notice that

$$D_q^z \neq \frac{1}{2}(D_q^x - iD_q^y), \quad D_q^w \neq \frac{1}{2}(D_q^u - iD_q^v).$$

The simplest set of double q -analytic functions is given by complex q -binomials

$$(z+iw)_q^n \equiv (z+iw)(z+iqw)(z+iq^2w)\dots(z+iq^{n-1}w) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{k(k-1)/2} i^k z^{n-k} w^k$$

satisfying

$$\frac{1}{2}(D_q^z + iD_q^w)(z + iw)_q^n = 0,$$

and

$$\frac{1}{2}(D_q^z - iD_q^w)(z + iw)_q^n = [n]_q(z + iw)_q^{n-1}.$$

Proof 4.0.5

$$\begin{aligned} & \frac{1}{2}(D_q^z - iD_q^w)(z + iw)_q^n \\ &= \frac{1}{2} \left(\sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{k(k-1)/2} i^k (D_q^z z^{n-k}) w^k - i \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{k(k-1)/2} i^k z^{n-k} \left(D_q^w w^k \right) \right) \\ &= \frac{1}{2} \left(\sum_{k=0}^{n-1} \begin{bmatrix} n \\ k \end{bmatrix}_q q^{k(k-1)/2} [n-k]_q z^{n-k-1} i^k w^k - i \sum_{k'=1}^n \begin{bmatrix} n \\ k' \end{bmatrix}_q q^{k'(k'-1)/2} i^{k'} z^{n-k'} \frac{w^{k'-1}}{q^{k'-1}} [k']_q \right) \\ &= \frac{1}{2} \left(\sum_{k=0}^{n-1} \begin{bmatrix} n \\ k \end{bmatrix}_q q^{k(k-1)/2} [n-k]_q z^{n-k-1} i^k w^k - i \sum_{k=0}^{n-1} \begin{bmatrix} n \\ k+1 \end{bmatrix}_q q^{k(k+1)/2} i^{k+1} z^{n-k-1} \frac{w^k}{q^k} [k+1]_q \right) \\ &= \frac{1}{2} \sum_{k=0}^{n-1} \left(\begin{bmatrix} n \\ k \end{bmatrix}_q q^{k(k-1)/2} [n-k]_q - i \begin{bmatrix} n \\ k+1 \end{bmatrix}_q q^{k(k+1)/2} i \frac{1}{q^k} [k+1]_q \right) z^{n-k-1} i^k w^k \\ &= \frac{1}{2} \sum_{k=0}^{n-1} 2 \left(\frac{[n]!}{[n-k-1]![k]!} q^{k(k-1)/2} \right) z^{n-k-1} i^k w^k \\ &= [n] \sum_{k=0}^{n-1} \begin{bmatrix} n-1 \\ k \end{bmatrix}_q q^{k(k-1)/2} z^{n-k-1} i^k w^k \\ &= [n](z + iw)_q^{n-1}. \end{aligned}$$

From above result follows that any convergent power series

$$f(z + iw)_q = \sum_{n=0}^{\infty} a_n (z + iw)_q^n$$

determines a double q -analytic function. Since our relation (44) shows expansion of double q -analytic q -binomials in terms of q -Hermite polynomials, it also gives expansion of any double q -analytic function in terms of the analytic polynomials.

Examples: For $n = 1$:

$$\begin{aligned} (z + iw)_q^1 &= z + iw = \frac{1}{[2]_q} \sum_{k=0}^1 \begin{bmatrix} 1 \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} H_{1-k}(z; q) H_k(qw; \frac{1}{q}) i^k \\ &= \frac{1}{[2]_q} \left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}_q H_1(z; q) H_0(qw; \frac{1}{q}) + \begin{bmatrix} 1 \\ 1 \end{bmatrix}_q H_0(z; q) H_1(qw; \frac{1}{q}) i \right) \end{aligned} \quad (53)$$

For $n = 2$:

$$\begin{aligned}
(z + iw)_q^2 &= (z + iw)(z + iqw) = z^2 + i[2]_q zw - qw^2 \\
&= \frac{1}{[2]_q^2} \sum_{k=0}^2 \begin{bmatrix} 2 \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} H_{2-k}(z; q) H_k(qw; \frac{1}{q}) i^k \\
&= \frac{1}{[2]_q^2} \left(\begin{bmatrix} 2 \\ 0 \end{bmatrix}_q H_2(z; q) H_0(qw; \frac{1}{q}) + \begin{bmatrix} 2 \\ 1 \end{bmatrix}_q H_1(z; q) H_1(qw; \frac{1}{q}) i + \begin{bmatrix} 2 \\ 2 \end{bmatrix}_q q H_0(z; q) H_2(qw; \frac{1}{q}) i^2 \right)
\end{aligned}$$

4.1 q -Holomorphic Laplacian

Another proof of identity (44) can be done by noticing that q -binomial $(z + iw)_q^n$ is double q -analytic function. Then we can use the following identity and complex q -Laplace equation.

Identity 4.1.1

$$e_q\left(-\frac{1}{[2]_q^2} \Delta_q\right)_q (z + iw)_q^n = (z + iw)_q^n \quad (54)$$

Proof 4.1.2 Since $(z + iw)_q^n$ is q -analytic function, it satisfies the q -Laplace equation

$$\Delta_q(z + iw)_q^n = 0$$

and

$$\begin{aligned}
e_q\left(-\frac{1}{[2]_q^2} \Delta_q\right)_q (z + iw)_q^n &= e_q\left(-\frac{1}{[2]_q^2} \left((D_q^z)^2 + (D_q^w)^2\right)\right)_q (z + iw)_q^n \\
&= \sum_{n=0}^{\infty} \frac{1}{[n]_q!} \left(-\frac{1}{[2]_q^2}\right)^n \left((D_q^z)^2 + (D_q^w)^2\right)_q^n (z + iw)_q^n \\
&= \sum_{n=0}^{\infty} \frac{1}{[n]_q!} \left(-\frac{1}{[2]_q^2}\right)^n (\Delta_q)_q^n (z + iw)_q^n, \quad (55)
\end{aligned}$$

where $(\Delta_q)_q^n = \Delta_q \cdot \Delta_q^{(1)} \cdot \Delta_q^{(2)} \cdot \dots \cdot \Delta_q^{(n-1)}$ and

$$\Delta_q = (D_q^z)^2 + (D_q^w)^2, \quad \Delta_q^{(1)} = (D_q^z)^2 + q(D_q^w)^2, \quad \dots, \quad \Delta_q^{(n-1)} = (D_q^z)^2 + q^{n-1}(D_q^w)^2.$$

Using the fact that $(\Delta_q)_q^m (z + iw)_q^n = 0, \forall m = 1, 2, \dots$, only the first term in expansion survives, then we get desired result.

Due to (47) we can factorize q -exponential operator function as

$$\begin{aligned}
&e_q\left(-\frac{1}{[2]_q^2} \Delta_q\right)_q (z + iw)_q^n \\
&= e_q\left(-\frac{1}{[2]_q^2} (D_q^z)^2\right)_q e_{\frac{1}{q}}\left(-\frac{1}{[2]_q^2} (D_q^w)^2\right)_q (z + iw)_q^n \\
&= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} i^k e_q\left(-\frac{1}{[2]_q^2} (D_q^z)^2\right)_q z^{n-k} e_{\frac{1}{q}}\left(-\frac{1}{[2]_q^2} (D_q^w)^2\right)_q w^k. \quad (56)
\end{aligned}$$

By using the generating function of q -Hermite Polynomials (45) we have the following identity:

$$H_n(x; q) = [2]_q^n e_q \left(-\frac{1}{[2]_q^2} (D_q^x)^2 \right) x^n, \quad (57)$$

which gives

$$e_q \left(-\frac{1}{[2]_q^2} (D_q^z)^2 \right) z^{n-k} = \frac{1}{[2]_q^{n-k}} H_{n-k}(z; q), \quad (58)$$

and

$$e_{\frac{1}{q}} \left(-\frac{1}{[2]_{\frac{1}{q}}^2} (D_{\frac{1}{q}}^w)^2 \right) w^k = \frac{1}{[2]_{\frac{1}{q}}^k q^k} H_k(qw; \frac{1}{q}), \quad (59)$$

where $D_{\frac{1}{q}}^{qw} = \frac{1}{q} D_{\frac{1}{q}}^w$. Substituting into (56), we get

$$e_q \left(-\frac{1}{[2]_q^2} \Delta_q \right)_q (z + iw)_q^n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} i^k \frac{1}{[2]_q^{n-k}} H_{n-k}(z; q) \frac{1}{[2]_{\frac{1}{q}}^k q^k} H_k(qw; \frac{1}{q}).$$

Then, according to identity (54) we obtain desired result

$$(z + iw)_q^n = \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} H_{n-k}(z; q) H_k(qw; \frac{1}{q}) i^k. \quad \diamond \quad (60)$$

As a particular case of our binomial formula, we can find q -Hermite binomial expansion for the q -analytic binomial $(x + iy)^n$ as well. If in (44) we replace $z \rightarrow x$ and $w \rightarrow y$, then we get

$$(x + iy)_q^n = \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} H_{n-k}(x; q) H_k(qy; \frac{1}{q}) i^k. \quad (61)$$

Since a q -analytic function is determined by power series in q -binomials [5], this formula allows us to get expansion of an arbitrary q -analytic function in terms of real q -Hermite polynomials.

5 q -Traveling Waves

As an application of q -binomials here we consider the q -analogue of traveling waves as a solution of q -wave equation.

5.1 Traveling Waves:

Real functions of two real variables $F(x, t) = F(x \pm ct)$ called the traveling waves, satisfy the following first order equations

$$\left(\frac{\partial}{\partial t} \mp c \frac{\partial}{\partial x} \right) F(x \pm ct) = 0.$$

It describes waves with fixed shape, prorogating with constant speed c in the left and in the right direction correspondingly. The general solution of the wave equation

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2},$$

then can be written as an arbitrary superposition of these traveling waves

$$u(x, t) = F(x + ct) + G(x - ct).$$

5.2 q-Traveling Waves

Direct extension of traveling waves to q-traveling waves is not possible. This happens due to the absence in q-calculus of the chain rule and as follows, impossibility to use moving frame as an argument of the wave function. Moreover, if we try in the Fourier harmonics $f(x, t) = e^{i(kx - \omega t)}$, replace exponential function by Jackson's q -exponential function $f(x, t) = e_q(i(kx - \omega t))$, then we find that it doesn't work due to the absence of factorization for q -exponential function $e_q(i(kx - \omega t)) \neq e_q(ikx) e_q(i\omega t)$.

This is why, here we propose another way. First we observe that q-binomials

$$(x \pm ct)_q^n = (x \pm ct)(x \pm qct) \dots (x \pm q^{n-1}ct) \quad (62)$$

for $n = 0, \pm 1, \pm 2, \dots$, satisfy the first order one-directional q -wave equations

$$\left(D_{\frac{1}{q}}^t \mp c D_q^x \right) (x \pm ct)_q^n = 0. \quad (63)$$

Then, the Laurent series expansion in terms of these q -binomials determines the q -analog of traveling waves

$$f(x \pm ct)_q = \sum_{n=-\infty}^{\infty} a_n (x \pm ct)_q^n.$$

Due to (63) the q -binomials (62) satisfy the q -wave equation

$$\left(\left(D_{\frac{1}{q}}^t \right)^2 - c^2 \left(D_q^x \right)^2 \right) u(x, t) = 0 \quad (64)$$

and the general solution of this equation is expressed in the form of q -traveling waves

$$u(x, t) = F(x + ct)_q + G(x - ct)_q \quad (65)$$

where

$$F(x + ct)_q = \sum_{n=-\infty}^{\infty} a_n (x + ct)_q^n$$

and

$$G(x - ct)_q = \sum_{n=-\infty}^{\infty} b_n (x - ct)_q^n.$$

This allows us to solve IVP for the q -wave equation

$$\left[\left(D_{\frac{1}{q}}^t \right)^2 - c^2 (D_q^x)^2 \right] u(x, t) = 0, \quad (66)$$

$$u(x, 0) = f(x), \quad (67)$$

$$D_{\frac{1}{q}}^t u(x, 0) = g(x), \quad (68)$$

where $-\infty < x < \infty$, in the D'Alembert form:

$$u(x, t) = \frac{f(x+ct)_q + f(x-ct)_q}{2} + \frac{1}{2c} \int_{(x-ct)_q}^{(x+ct)_q} g(x') d_q x', \quad (69)$$

where the Jackson integral is

$$\begin{aligned} \int_{(x-ct)_q}^{(x+ct)_q} g(x') d_q x' &= (1-q)(x+ct) \sum_{j=0}^{\infty} q^j g(q^j(x+ct))_q \\ &\quad - (1-q)(x-ct) \sum_{j=0}^{\infty} q^j g(q^j(x-ct))_q. \end{aligned} \quad (70)$$

If the initial velocity is zero, $g(x) = 0$, the formula reduces to

$$u(x, t) = \frac{1}{2} (f(x+ct)_q + f(x-ct)_q). \quad (71)$$

It should be noted here that q -traveling wave is not traveling wave in the standard sense and it is not preserving shape during evolution. It can be seen from simple observation. The traveling wave polynomial $(x-ct)_q^n = (x-ct)(x-qct)(x-q^2ct)\dots(x-q^{n-1}ct)$ includes the set of moving frames (as zeros of this polynomial) with re-scaled set of speeds $(c, qc, q^2c, \dots, q^{n-1}c)$. It means that zeros of this polynomial are moving with different speeds and therefore the shape of polynomial wave is not preserving. Only in the linear case and in the case $q = 1$, when speeds of all frames coincide, we are getting standard traveling wave.

5.3 EXAMPLES

In this section we are going to illustrate our results by several explicit solutions.

Example 1: We consider I.V.P. for the q -wave equation (66) with initial functions

$$\begin{aligned} u(x, 0) &= x^2, \\ D_{\frac{1}{q}}^t u(x, 0) &= 0. \end{aligned} \quad (72)$$

Then the solution of the given I.V.P. for q -wave equation in D'Alembert form is found as

$$u(x, t) = x^2 + qc^2t^2. \quad (73)$$

When $q = 1$, it reduces to well-known one as superposition of two traveling wave parabolas $(x \pm ct)^2$ moving to the right and to the left with speed c . Geometrically, the meaning of q is the acceleration of our parabolas in vertical direction.

Example 2: The q -traveling wave

$$\begin{aligned} u(x, t) = (x - ct)_q^2 &= (x - ct)(x - qct) \\ &= \left(x - \frac{[2]_q}{2}ct\right)^2 - \frac{(q-1)^2}{4}c^2t^2 \end{aligned} \quad (74)$$

gives solution of I.V.P. for the q -wave equation (66) with initial functions

$$\begin{aligned} u(x, 0) &= x^2 \\ D_{\frac{1}{q}}^t u(x, 0) &= -[2]_q cx. \end{aligned} \quad (75)$$

If $q = 1$ in this solution we have two degenerate zeros moving with the same speed c . In the case $q \neq 1$, two zeros are moving with different speeds c and qc . It means that, the distance between zeros is growing linearly with time as $(q-1)ct$. The solution is the parabola, moving in vertical direction with acceleration $\frac{(q-1)^2}{4}c^2$, and in horizontal direction with constant speed $\frac{[2]_q}{2}c$. The area under the curve between moving zeros $x = ct$ and $x = qct$

$$\int_{ct}^{qct} (x - ct)_q^2 dx = -\frac{(q-1)^3 c^3}{6} t^3$$

is changing according to time as t^3 .

For more general initial function $f(x) = x^n$, $n = 2, 3, \dots$ we get q -traveling wave

$$u(x, t) = (x - ct)_q^n = (x - ct)(x - qct) \dots (x - q^{n-1}ct)$$

with n -zeros moving with speeds $c, qc, \dots, q^{n-1}c$. The distance between two zeros is growing as $(q^m - q^n)ct$, and the shape of wave is changing. In parabolic case with $n = 2$, the shape of curve is not changing, but moving in horizontal direction with constant speed, and in vertical direction with constant acceleration. In contrast to this, for $n > 2$, the motion of zeros with different speeds changes the shape of the wave, and it can not be reduced to simple translation and acceleration.

Example 3: Given I.V.P. for the q -wave equation (66) with initial functions as q -trigonometric functions [3]

$$\begin{aligned} u(x, 0) &= \cos_q x, \\ D_{\frac{1}{q}}^t u(x, 0) &= \sin_q x. \end{aligned} \quad (76)$$

By using the D'Alembert form (69), after q -integration, we get

$$\begin{aligned} u(x, t) &= \frac{1}{2} [\cos_q(x + ct)_q + \cos_q(x - ct)_q] + \frac{1}{2c} \int_{(x-ct)_q}^{(x+ct)_q} \sin_q(x') d_q x' \\ &= \frac{1}{2} \left[\left(1 + \frac{1}{c}\right) \cos_q(x - ct)_q + \left(1 - \frac{1}{c}\right) \cos_q(x + ct)_q \right]. \end{aligned} \quad (77)$$

Example 4: q -Gaussian Traveling Wave For initial function in Gaussian form: $u(x, 0) = e^{-x^2}$ in the standard case $q = 1$ we have the Gaussian traveling wave $u(x, t) = e^{-(x-ct)^2}$. For the q -traveling wave with Gaussian initial condition $u(x, 0) = e^{-x^2}$, we have the q -traveling wave

$$u(x, t) = \left(e^{-(x-ct)^2} \right)_q \equiv \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} (x-ct)_q^{2n}.$$

5.4 q -Traveling Waves in terms of q -Hermite Polynomials

Identity(25) allows us to rewrite the traveling wave binomial in terms of Hermite polynomials as

$$(x+ct)^n = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} i^k H_{n-k}(x) H_k(-ict).$$

Its q -analogue for q -traveling wave binomial follows from (44)

$$(x+ct)_q^n = \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q i^k q^{\frac{k(k-1)}{2}} H_{n-k}(x; q) H_k(-iqct, \frac{1}{q}).$$

Then, the general solution of q -wave equation (64) can be expressed in the form of q -Hermite polynomials

$$u(x, t) = F(x+ct)_q + G(x-ct)_q, \quad (78)$$

where

$$\begin{aligned} F(x+ct)_q &= \sum_{n=-\infty}^{\infty} a_n (x+ct)_q^n = \sum_{n=-\infty}^{\infty} a_n \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q i^k q^{\frac{k(k-1)}{2}} H_{n-k}(x; q) H_k(-iqct, \frac{1}{q}), \\ G(x-ct)_q &= \sum_{n=-\infty}^{\infty} a_n (x-ct)_q^n = \sum_{n=-\infty}^{\infty} a_n \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q i^k q^{\frac{k(k-1)}{2}} H_{n-k}(x; q) H_k(iqct, \frac{1}{q}). \end{aligned}$$

It is instructive to prove the q -traveling wave solution

$$\left(D_{\frac{1}{q}}^t - cD_q^x \right) (x+ct)_q^n = 0$$

by using q -Hermite binomial. We have

$$\begin{aligned} (D_{\frac{1}{q}}^t - cD_q^x)(x+ct)_q^n &= \left(D_{\frac{1}{q}}^t - cD_q^x \right) \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q i^k q^{\frac{k(k-1)}{2}} H_{n-k}(x; q) H_k(-iqct, \frac{1}{q}) \\ &= \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q i^k q^{\frac{k(k-1)}{2}} H_{n-k}(x; q) D_{\frac{1}{q}}^t H_k(-iqct, \frac{1}{q}) \\ &\quad - c \frac{1}{[2]_q^n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q i^k q^{\frac{k(k-1)}{2}} D_q^x H_{n-k}(x; q) H_k(-iqct, \frac{1}{q}) \end{aligned}$$

By recursion formula for q -Hermite polynomials

$$D_q^x H_n(x; q) = [2]_q [n]_q H_{n-1}(x; q)$$

we get

$$\begin{aligned} (D_{\frac{1}{q}}^t - cD_q^x)(x + ct)_q^n &= \frac{1}{[2]_q^n} \sum_{k'=1}^n \begin{bmatrix} n \\ k' \end{bmatrix}_q i^{k'} q^{\frac{k'(k'-1)}{2}} H_{n-k'}(x; q) [2]_{\frac{1}{q}} [k']_{\frac{1}{q}} H_{k'-1}(-iqct, \frac{1}{q})(-iqc) \\ &\quad - c \frac{1}{[2]_q^n} \sum_{k=0}^{n-1} \begin{bmatrix} n \\ k \end{bmatrix}_q i^k q^{\frac{k(k-1)}{2}} [2]_q [n-k]_q H_{n-k-1}(x; q) H_k(-iqct, \frac{1}{q}) \\ &= \frac{1}{[2]_q^n} \sum_{k=0}^{n-1} \begin{bmatrix} n \\ k+1 \end{bmatrix}_q i^{k+1} q^{\frac{k(k+1)}{2}} H_{n-k-1}(x; q) [2]_{\frac{1}{q}} [k+1]_{\frac{1}{q}} H_k(-iqct, \frac{1}{q})(-iqc) \\ &\quad - c \frac{1}{[2]_q^n} \sum_{k=0}^{n-1} \begin{bmatrix} n \\ k \end{bmatrix}_q i^k q^{\frac{k(k-1)}{2}} [2]_q [n-k]_q H_{n-k-1}(x; q) H_k(-iqct, \frac{1}{q}) \\ &= ci^k \frac{1}{[2]_q^n} \sum_{k=0}^{n-1} \left(\begin{bmatrix} n \\ k+1 \end{bmatrix}_q i q^{\frac{k(k+1)}{2}} [2]_{\frac{1}{q}} [k+1]_{\frac{1}{q}} - \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} [2]_q [n-k]_q \right) H_{n-k-1}(x; q) H_k(-iqct, \frac{1}{q}) \\ &= 0. \end{aligned}$$

The expression in parenthesis is zero due to q -combinatorial formula and $[n]_{\frac{1}{q}} = \frac{[n]_q}{q^{n-1}}$.

Acknowledgments

This work was supported by Izmir Institute of Technology. One of the authors (S. Nalci) was partially supported by TUBITAK scholarship for graduate students.

References

- [1] A.Perelomov, Generalized Coherent states and their applications, Springer-Verlag,1986.
- [2] E. G. Floratos, The many-body problem for q -oscillators J. Phys. A: Math. Gen. 24,1991.
- [3] V. Kac and P. Cheung, Quantum Calculus, Springer, New York, 2002.
- [4] S.Nalci and O.K.Pashaev, q -analog of shock soliton solution, J. Phys. A: Math. Theor. 43, 445205, 2010.
- [5] O.K. Pashaev and S.Nalci, q -analytic functions, fractals and generalized analytic functions, J. Phys. A: Math. Theor. 47,045204,2014.

- [6] G.Dattoli et all ,Theory of generalized Hermite polynomials, Computer Math. Applic. Vol:28,N0:4, pp.71-83, 1994.